

January 8, 2004

Residual Autocorrelation Testing for Vector Error Correction Models ¹

Ralf Brüggemann

European University Institute, Florence and Humboldt University, Berlin

Helmut Lütkepohl

European University Institute, Florence and Humboldt University, Berlin

Pentti Saikkonen

University of Helsinki

Address for correspondence: Helmut Lütkepohl, Economics Department, European University Institute, Via della Piazzuola 43, I-50133 Firenze, ITALY
email: helmut.luetkepohl@iue.it

Abstract

In applied time series analysis, checking for autocorrelation in a fitted model is a routine diagnostic tool. Therefore it is useful to know the asymptotic and small sample properties of the standard tests for the case when some of the variables are cointegrated. The properties of residual autocorrelations of vector error correction models (VECMs) and tests for residual autocorrelation are derived. In particular, the asymptotic distributions of Lagrange multiplier (LM) and portmanteau tests are given. Monte Carlo simulations show that the LM tests have advantages if autocorrelation of small order is tested whereas portmanteau tests are preferable for higher order residual autocorrelation. Their critical values have to be adjusted for the cointegration rank of the system, however.

JEL classification: C32

Keywords: Cointegration, dynamic econometric models, vector autoregressions, vector error correction models, residual autocorrelation

¹The third author thanks the Yrjö Jahnsson Foundation, the Academy of Finland and the Alexander von Humboldt Foundation under a Humboldt research award for financial support. Part of this research was done while he was visiting the European University Institute in Florence.

1 Introduction

When a model for the data generation process (DGP) of a time series or a set of time series has been constructed, it is common to perform checks of the model adequacy. Tests for residual autocorrelation (AC) are prominent tools for this task. A number of such tests are in routine use for both stationary as well as nonstationary processes with integrated or cointegrated variables. Well-known examples are portmanteau and Lagrange multiplier (LM) tests for residual AC. They are, for example, available in commercial econometric software such as EViews and PcGive. The theoretical properties of these tests are well explored for stationary DGPs (see, e.g., Ahn (1988), Hosking (1980, 1981a, 1981b), Li & McLeod (1981) or Lütkepohl (1991) for a textbook exposition with more references and Edgerton & Shukur (1999) for a large scale small sample comparison of various tests). An explicit treatment of the properties of residual ACs of vector error correction models (VECMs) for cointegrated variables does not seem to be available, however. This may also be the reason why no critical values or p -values are presented in PcGive for some AC tests for VECMs. On the other hand, p -values based on the asymptotic distributions of portmanteau and LM tests for AC are provided in EViews 4 although a theoretical justification seems to be missing. In fact, it will be seen from our results that the degrees of freedom of the portmanteau tests should be adjusted in the presence of cointegration.

In this study it will be assumed that all variables are stationary or integrated of order one (I(1)) and the DGP is a VECM with cointegration rank r . The asymptotic distribution of the residual ACs is derived for a model estimated by reduced rank (RR) regression or some similar method. The crucial condition here is that the cointegration matrix is estimated superconsistently. In that case it turns out that the residual ACs have the same asymptotic distribution as if the cointegration relations were known. There may also be restrictions on the cointegration or short-run parameters.

Because standard tests for residual AC are based on the estimated autocovariances or ACs, the asymptotic properties of the tests follow from those of these quantities. In particular, we consider LM tests and portmanteau tests in more detail. While the former tests have the same asymptotic distribution as in the stationary case, the situation is different for the portmanteau tests. For the latter tests the degrees of freedom have to be adjusted to account for the cointegration rank of the system. Thus, the practice used in EViews 4,

for example, to subtract only the number of parameters associated with differenced lagged values from the number of autocorrelations included in the statistic when determining the degrees of freedom, has no asymptotic justification.

We also analyze the small sample properties of the tests in a Monte Carlo study. Not surprisingly, some of the tests have small sample properties for which the asymptotic theory is a poor guide. Similar results have also been found for stationary processes (Edgerton & Shukur (1999) and Doornik (1996)). It turns out, however, that for low-dimensional systems and moderately large samples, the usual tests perform reasonably well if the cointegration properties of the process are taken into account properly.

The study is organized as follows. In the next section the model setup is presented and the asymptotic properties of the residual autocovariances and ACs are discussed in Section 3. Tests for residual AC are considered in Section 4. The results of a Monte Carlo study are presented in Section 5 and concluding remarks follow in Section 6. Proofs are presented in the Appendix.

The following general notation will be used. The differencing and lag operators are denoted by Δ and L , respectively. Convergence in distribution is signified by \xrightarrow{d} and \log denotes the natural logarithm. $O_p(\cdot)$ and $o_p(\cdot)$ are the usual symbols for boundedness in probability. Independently, identically distributed is abbreviated by iid. The trace, determinant and rank of the matrix A are denoted by $\text{tr}(A)$, $\det(A)$ and $\text{rk}(A)$, respectively. The symbol vec is used for the column vectorization operator. If A is an $(n \times m)$ matrix of full column rank ($n > m$), we denote an orthogonal complement by A_\perp . The orthogonal complement of a nonsingular square matrix is zero and the orthogonal complement of a zero matrix is an identity matrix of suitable dimension. An $(n \times n)$ identity matrix is denoted by I_n . DGP, ML, LS, GLS, RR, LR and LM are used to abbreviate data generation process, maximum likelihood, least squares, generalized least squares, reduced rank, likelihood ratio and Lagrange multiplier, respectively. VAR and VECM stand for vector autoregression and vector error correction model, respectively. A sum is defined to be zero if the lower bound of the summation index exceeds the upper bound.

2 The Model Setup

Suppose $y_t = (y_{1t}, \dots, y_{Kt})'$ is the K -dimensional vector of observable I(1) time series variables with $r < K$ cointegration relations. They are assumed to be generated by a DGP which can be written in VECM form

$$\Delta y_t = \nu + \alpha(\beta' y_{t-1} - \tau(t-1)) + \Gamma_1 \Delta y_{t-1} + \dots + \Gamma_{p-1} \Delta y_{t-p+1} + u_t, \quad (2.1)$$

where α, β are $(K \times r)$ matrices with rank r , ν is a $(K \times 1)$ vector of constants, τ is an r -dimensional vector of trend slope parameters and $\Gamma_1, \dots, \Gamma_{p-1}$ are $(K \times K)$ short-run parameter matrices. The parameters are such that the DGP is I(1). The error process u_t is assumed to be iid with mean zero and nonsingular covariance matrix Ω , that is, $u_t \sim \text{iid}(0, \Omega)$.

It follows from Granger's representation theorem that

$$y_t = \mu_0 + \mu_1 t + x_t, \quad t = 1, 2, \dots, \quad (2.2)$$

where μ_0 and μ_1 are functions of the parameters in (2.1) and

$$x_t = \mathcal{C} \sum_{j=1}^t u_j + \mathcal{C}(L)u_t + P_{\beta_{\perp}} x_0 \quad (2.3)$$

with $\mathcal{C} = \beta_{\perp}(\alpha'_{\perp}(I - \sum_{j=1}^{p-1} \Gamma_j)\beta_{\perp})^{-1}\alpha'_{\perp}$, x_0 contains initial values, $P_{\beta_{\perp}} = \beta_{\perp}(\beta'_{\perp}\beta_{\perp})^{-1}\beta'_{\perp}$ and $\mathcal{C}(L)$ is a geometrically convergent power series in the lag operator L . The process x_t can also be written as

$$\Delta x_t = \alpha\beta' x_{t-1} + \Gamma_1 \Delta x_{t-1} + \dots + \Gamma_{p-1} \Delta x_{t-p+1} + u_t. \quad (2.4)$$

It is assumed that the parameters of the model (2.1) are estimated by RR regression (Gaussian ML estimation) or some other asymptotically equivalent method which, for example, allows for parameter restrictions. The estimators are indicated by a hat. Once estimators of β and τ are available, the estimators $\hat{\nu}$, $\hat{\alpha}$ and $\hat{\Gamma}_j$ ($j = 1, \dots, p-1$) are supposed to be obtained from (2.1) by LS or GLS with β and τ replaced by $\hat{\beta}$ and $\hat{\tau}$, respectively. For example, if the latter estimators are obtained by RR regression, then $\hat{\nu}$, $\hat{\alpha}$ and $\hat{\Gamma}_j$ are Gaussian ML estimators.

In the following section the asymptotic properties of the residual autocovariances and ACs will be considered.

3 Residual Autocovariances and Autocorrelations

We consider the residual autocovariances

$$\hat{C}_j = \frac{1}{T} \sum_{t=j+1}^T \hat{u}_t \hat{u}'_{t-j}, \quad j = 1, \dots, h, \quad (3.1)$$

where

$$\hat{u}_t = \Delta y_t - \hat{\nu} - \hat{\alpha}(\hat{\beta}' y_{t-1} - \hat{\tau}(t-1)) - \hat{\Gamma}_1 \Delta y_{t-1} - \dots - \hat{\Gamma}_{p-1} \Delta y_{t-p+1}.$$

Let $\hat{C} = [\hat{C}_1, \dots, \hat{C}_h]$ and $\hat{c} = \text{vec}(\hat{C})$. Under general conditions, this quantity has an asymptotic normal distribution. More precisely, we have the following result.

Proposition 1.

Suppose the assumptions of Section 2 hold. Let $U_t = [u'_{t-1}, \dots, u'_{t-h}]'$ and $X_t = [(\beta' x_{t-1})', \Delta x'_{t-1}, \dots, \Delta x'_{t-p+1}]'$ and define

$$\Sigma_{XU} = \text{plim } T^{-1} \sum_{t=1}^T X_t U_t' = \Sigma'_{UX}, \quad \Sigma_{XX} = \text{plim } T^{-1} \sum_{t=1}^T X_t X_t'. \quad (3.2)$$

Then,

$$T^{1/2} \hat{c} \xrightarrow{d} N(0, \Sigma_{U \cdot X} \otimes \Omega), \quad (3.3)$$

where $\Sigma_{U \cdot X} = \Sigma_{UU} - \Sigma_{UX} \Sigma_{XX}^{-1} \Sigma_{XU}$ with $\Sigma_{UU} = I_h \otimes \Omega$. □

The probability limits in (3.2) exist due to a weak law of large numbers because X_t is a linear process with geometrically decaying coefficients by (2.3) (see Johansen (1995, Appendix B.7)). The limiting distribution in (3.3) is analogous to the stationary VAR case (see, e.g., Ahn (1988) or Lütkepohl (1991)). It is the same that would be obtained if the cointegration matrix were known and the deterministic terms in the process were zero. This fact is used in proving the proposition. To see this, let μ_{0o} and μ_{1o} denote the true values of the parameters μ_0 and μ_1 , respectively, and make the transformation $y_t \rightarrow x_t = y_t - \mu_{0o} - \mu_{1o}t$ to obtain

$$\Delta x_t = \nu^{(0)} + \alpha(\beta' x_{t-1} - \tau^{(0)}(t-1)) + \Gamma_1 \Delta x_{t-1} + \dots + \Gamma_{p-1} \Delta x_{t-p+1} + u_t, \quad (3.4)$$

where $\nu^{(0)} = \nu + \alpha\beta'\mu_{0o} - (I_K - \sum_{j=1}^{p-1} \Gamma_j)\mu_{1o}$ and $\tau^{(0)} = \tau - \beta'\mu_{1o}$. It can be seen by expressing ν and τ in terms of μ_0 and μ_1 that the true values of the parameters $\nu^{(0)}$ and $\tau^{(0)}$

are zero and their estimators are $\hat{\nu}^{(0)} = \hat{\nu} + \hat{\alpha}\hat{\beta}'\mu_{0o} - (I_K - \sum_{j=1}^{p-1}\hat{\Gamma}_j)\mu_{1o}$ and $\hat{\tau}^{(0)} = \hat{\tau} - \hat{\beta}'\mu_{1o}$, respectively. Thus, the transformation from model (2.1) to (3.4) leaves the error term u_t and the estimated residuals \hat{u}_t unaltered and therefore the transformed model can be used to obtain the limiting distribution of the residual autocovariances \hat{C}_j . Doing so requires suitable assumptions for the employed estimators of the parameters in (2.1). Because the proofs are based on the transformed model (3.4), it is convenient to present these assumptions by using estimators of the parameters $\nu^{(0)}$ and $\tau^{(0)}$ instead of the original deterministic parameters ν and τ .

The estimators of β and τ are such that the following conditions hold:

$$\hat{\beta} = \beta + O_p(T^{-1}) \quad \text{and} \quad \hat{\tau}^{(0)} = O_p(T^{-3/2}). \quad (3.5)$$

It is well-known that these results hold if β and τ are estimated by RR regression and the estimator of β is normalized or identified in some appropriate way (see, e.g., Johansen (1995)). Any normalization of the type $\xi'\beta = I_r$ with ξ a $(K \times r)$ matrix of full column rank can be employed. Because the convergence result and, hence, the normalization is used for theoretical derivations only, ξ need not be known or specified and it does not have to be feasible in practice. For example, $\xi = \beta(\beta'\beta)^{-1}$ may be used.

As already mentioned in Section 2, the parameters ν , α and Γ_j ($j = 1, \dots, p-1$) are supposed to be estimated from (2.1) by LS or GLS with β and τ replaced by $\hat{\beta}$ and $\hat{\tau}$, respectively. It is well-known that we have

$$\hat{\nu}^{(0)} = O_p(T^{-1/2}), \quad \hat{\alpha} = \alpha + O_p(T^{-1/2}), \quad \hat{\Gamma}_j = \Gamma_j + O_p(T^{-1/2}) \quad \text{for } j = 1, \dots, p-1 \quad (3.6)$$

(see, e.g., Johansen (1995)). Notice that these results also hold, for example, if there are restrictions on the short-run parameters.

We shall show next that, under the foregoing assumptions, the limiting distribution in (3.3) is the same one would obtain from model (2.4) with known cointegrating vectors. Therefore suppose that the model (2.4) is estimated in the same way as model (2.1) except that the cointegration matrix β is assumed known. Denoting the estimated residuals by \tilde{u}_t and the corresponding residual autocovariances by \tilde{C}_j ($j = 1, \dots, h$), the following lemma is obtained. A proof is given in the Appendix.

Lemma 1.

$$\tilde{C}_j - \hat{C}_j = O_p(T^{-1}), \quad j = 1, \dots, h. \quad (3.7)$$

□

A proof of Proposition 1 based on this lemma is also given in the Appendix. It follows immediately from Proposition 1 that an analogous result also holds for the residual ACs. Notice that the AC matrix corresponding to \hat{C}_j is $\hat{R}_j = \hat{S}^{-1}\hat{C}_j\hat{S}^{-1}$, where \hat{S} is a diagonal matrix with the square roots of the diagonal elements of $\hat{\Omega} = T^{-1}\sum_{t=1}^T \hat{u}_t\hat{u}_t'$ on the diagonal. Moreover, let S be the corresponding diagonal matrix with the square roots of the diagonal elements of Ω on the diagonal and let $R_0 = S^{-1}\Omega S^{-1}$ be the correlation matrix corresponding to Ω . With this notation we can state the following corollary of Proposition 1.

Corollary 1.

Let $\hat{R} = [\hat{R}_1, \dots, \hat{R}_h]$ and $\hat{\mathbf{r}} = \text{vec}(\hat{R}) = (I_h \otimes \hat{S}^{-1} \otimes \hat{S}^{-1})\hat{\mathbf{c}}$. Then, under the conditions of Proposition 1,

$$T^{1/2}\hat{\mathbf{r}} \xrightarrow{d} N(0, [(I_h \otimes S^{-1})\Sigma_{U.X}(I_h \otimes S^{-1})] \otimes R_0). \quad (3.8)$$

□

This result can be used to set up confidence intervals around zero for the residual ACs of an estimated VECM in the same way as for stationary processes (see, e.g., Lütkepohl (1991)). As in the latter case, the asymptotic standard errors are smaller than the crude $1/\sqrt{T}$ approximations which are sometimes used as a rough check for residual AC. A couple of extensions may be worth noting.

Extension 1. The results also hold if there is no deterministic trend term in the model. In other words, $\tau = 0$ is known a priori. Moreover, seasonal dummies may be added to the model. They do not have an effect on the asymptotic distributions of the residual autocovariances and ACs.

Extension 2. There may be linear or smooth nonlinear restrictions on the parameters α

and Γ_j ($j = 1, \dots, p-1$). Zero restrictions are, of course, a leading case of such restrictions. Suppose that $\alpha = \alpha(\theta)$ and $\Gamma_j = \Gamma_j(\theta)$ ($j = 1, \dots, p-1$) with θ an underlying structural parameter. Suppose the estimator of θ is consistent and asymptotically normal. Then the previous results continue to hold with the usual adjustments to the asymptotic covariance matrix of the parameter estimators and, hence of $\Sigma_{U.X}$ (see Ahn (1988) or Lütkepohl (1991)), if the corresponding estimators of α and Γ_j satisfy (3.6) and the estimators of β and τ maintain their convergence rates and are asymptotically independent of the estimator of θ . It has been discussed by Johansen (1991, Appendix) and Reinsel & Ahn (1992) that the required asymptotic estimation theory holds for restricted versions of model (2.1). One restriction of particular importance is obtained by confining the intercept term to the cointegration relations. This constraint may be imposed if there are no linear deterministic trends in the variables. It is considered by Saikkonen (2001a, b).

Extension 3. The same asymptotic distributions are also obtained for the residual autocovariances and ACs if the model (2.1) is estimated without the cointegration restriction, that is, the cointegration rank r is ignored and no restriction is placed on the error correction term. Instead of (2.1) we then consider

$$\Delta y_t = \nu + \nu_1(t-1) + \Pi y_{t-1} + \Gamma_1 \Delta y_{t-1} + \dots + \Gamma_{p-1} \Delta y_{t-p+1} + u_t, \quad (3.9)$$

where $\nu_1 = -\alpha\tau$ and $\Pi = \alpha\beta'$. The parameters are assumed to be estimated by LS and the resulting estimators are again indicated by a hat. The error term u_t and the LS residuals \hat{u}_t are again invariant to subtracting the deterministic term so that the corresponding true parameter values may be assumed to be zero and $y_t = x_t$ can be assumed in (3.9). Using further the transformation

$$\Pi x_{t-1} = \alpha\beta' [\xi_o : \beta_{o\perp}] \begin{bmatrix} \beta'_o \\ \xi'_{o\perp} \end{bmatrix} x_{t-1} = a(\beta'_o x_{t-1}) + b(\xi'_{o\perp} x_{t-1}),$$

where $a = \alpha\beta'_o \xi_o$, $b = \alpha\beta' \beta_{o\perp}$ and ‘ o ’ refers to true parameter values, the limiting distribution of the residual autocovariances can be derived by replacing (3.9) by

$$\Delta x_t = \nu^{(0)} + \nu_1^{(0)}(t-1) + av_{0t} + bv_{1t} + \Gamma_1 \Delta x_{t-1} + \dots + \Gamma_{p-1} \Delta x_{t-p+1} + u_t, \quad (3.10)$$

where $v_{0t} = \beta'_o x_{t-1}$, $v_{1t} = \xi'_{o\perp} x_{t-1}$, $\nu^{(0)}$ is as defined before and $\nu_1^{(0)} = -\alpha\tau^{(0)}$ so that the true values of the parameters $\nu^{(0)}$, $\nu_1^{(0)}$ and b are zero. The processes Δx_t and v_{0t} are stationary

and have zero mean whereas v_{1t} is I(1) and not cointegrated. The LS residuals \hat{u}_t from (3.9) can be considered as LS residuals from (3.10). In the same way as in the proof of Lemma 1, we can use well-known limit results of stationary and I(1) processes and obtain an analog of Lemma 1 with \hat{C}_j residual autocovariance matrices based on (3.10) and \tilde{C}_j LS residual autocovariance matrices obtained from (3.10) with the constraints $\nu = 0$, $\nu_1 = 0$ and $b = 0$ (see (A.2) in the Appendix). Thus, it suffices to obtain the limiting distribution of \tilde{C}_j ($j = 1, \dots, h$) and Proposition 1 can be applied directly.

4 Testing for Residual Autocorrelation

Two types of tests for residual AC are quite popular in applied work, Breusch-Godfrey LM tests and portmanteau tests. They are both based on statistics of the form

$$Q = T\hat{\mathbf{c}}'\hat{\Sigma}^{-1}\hat{\mathbf{c}}. \quad (4.1)$$

where $\hat{\Sigma}$ is a suitable scaling matrix. In other words, they are based on the residual autocovariances. The choice of scaling matrix $\hat{\Sigma}$ determines the type of test statistic and its asymptotic distribution under the null hypothesis of no residual AC. For the Breusch-Godfrey LM statistic, $\hat{\Sigma}$ is an estimator of the covariance matrix of the limiting distribution in (3.3) of Proposition 1, whereas the portmanteau statistic uses an estimator of $I_h \otimes \Omega \otimes \Omega$ instead. We will consider both types of tests in turn.

4.1 Breusch-Godfrey Test

The Breusch-Godfrey test can be viewed as considering a VAR(h) model

$$u_t = B_1 u_{t-1} + \dots + B_h u_{t-h} + \varepsilon_t \quad (4.2)$$

for the error terms and testing the pair of hypotheses

$$H_0 : B_1 = \dots = B_h = 0 \quad \text{vs.} \quad H_1 : B_1 \neq 0 \text{ or } \dots \text{ or } B_h \neq 0. \quad (4.3)$$

For this purpose the auxiliary model

$$\begin{aligned} \hat{u}_t &= B_1 \hat{u}_{t-1} + \dots + B_h \hat{u}_{t-h} + \alpha \hat{\beta}^{*'} y_{t-1}^* + \Gamma_1 \Delta y_{t-1} + \dots + \Gamma_{p-1} \Delta y_{t-p+1} \\ &\quad + \nu + (y'_{t-1} \xi_{\perp} \otimes \hat{\alpha}) \phi_{11} - \hat{\alpha}(t-1) \phi_{21} + e_t \\ &= (\hat{U}'_t \otimes I_K) \gamma + (\hat{Z}'_t \otimes I_K) \phi + \hat{Z}'_{1t} \phi_1 + e_t, \quad t = 1, \dots, T, \end{aligned} \quad (4.4)$$

may be used with $\hat{\beta}^{*'} = [\hat{\beta}' : \hat{\tau}]$, $y_{t-1}^{*'} = [y'_{t-1} : t-1]$, $\hat{U}'_t = [\hat{u}'_{t-1}, \dots, \hat{u}'_{t-h}]$ ($\hat{u}_t = 0$ for $t \leq 0$), $\gamma = \text{vec}[B_1, \dots, B_h]$, $\hat{Z}'_t = [(\hat{\beta}^{*'} y_{t-1}^{*'})', \Delta y'_{t-1}, \dots, \Delta y'_{t-p+1}]$, $\phi = \text{vec}[\alpha, \Gamma_1, \dots, \Gamma_{p-1}]$, $\hat{Z}'_{1t} = [I_K : y'_{t-1} \xi_{\perp} \otimes \hat{\alpha} : -\hat{\alpha}(t-1)]$ and $\phi'_1 = [\nu', \phi'_{11}, \phi'_{21}]$. The terms in \hat{Z}'_{1t} are related to the scores of ν , β and τ . Allowing for ν simply means including an intercept term in (4.4) which is again denoted by ν for simplicity. Regarding β , the situation depends on a possible normalization. If a normalization $\xi' \beta = I_r$ with ξ known is used, the natural additional regressor is $y'_{t-1} \xi_{\perp} \otimes \hat{\alpha}$. If a particular normalization is not used, the choice $\xi_{\perp} = \hat{\beta}_{\perp}$ can be considered. The terms $y'_{t-1} \xi_{\perp} \otimes \hat{\alpha}$ and $-\hat{\alpha}(t-1)$ may in fact be deleted from the auxiliary model (see Remark 1 below).

The Breusch-Godfrey test statistic, say Q_{BG}^* , is a standard LM test statistic for the null hypothesis $\gamma = 0$ in (4.4),

$$Q_{BG}^* = T \hat{\gamma}' (\hat{\Sigma}^{\gamma\gamma})^{-1} \hat{\gamma} = T \hat{\mathbf{c}}' (I_{hK} \otimes \hat{\Omega}^{-1}) \hat{\Sigma}^{\gamma\gamma} (I_{hK} \otimes \hat{\Omega}^{-1}) \hat{\mathbf{c}}, \quad (4.5)$$

where $\hat{\gamma}$ is the GLS estimator of γ and $\hat{\Sigma}^{\gamma\gamma}$ is the part of

$$\left(\begin{array}{c} T^{-1} \sum_{t=1}^T \begin{bmatrix} \hat{U}_t \otimes I_K \\ \hat{Z}_t \otimes I_K \\ \hat{Z}_{1t} \end{bmatrix} \hat{\Omega}^{-1} [\hat{U}'_t \otimes I_K : \hat{Z}'_t \otimes I_K : \hat{Z}'_{1t}] \end{array} \right)^{-1}$$

corresponding to γ . That is,

$$\hat{\Sigma}^{\gamma\gamma} = \left(\Sigma_{\hat{U}\hat{U}} - \Sigma_{\hat{U}\hat{Z}} \Sigma_{\hat{Z}\hat{Z}}^{-1} \Sigma'_{\hat{U}\hat{Z}} \right)^{-1}$$

with

$$\Sigma_{\hat{U}\hat{U}} = T^{-1} \sum_{t=1}^T (\hat{U}_t \otimes I_K) \hat{\Omega}^{-1} (\hat{U}_t \otimes I_K)',$$

$$\Sigma_{\hat{U}\hat{Z}} = T^{-1} \sum_{t=1}^T (\hat{U}_t \otimes I_K) \hat{\Omega}^{-1} [\hat{Z}'_t \otimes I_K : \hat{Z}'_{1t}]$$

and

$$\Sigma_{\hat{Z}\hat{Z}} = T^{-1} \sum_{t=1}^T \begin{bmatrix} \hat{Z}_t \otimes I_K \\ \hat{Z}_{1t} \end{bmatrix} \hat{\Omega}^{-1} [\hat{Z}'_t \otimes I_K : \hat{Z}'_{1t}].$$

Here $\hat{\Omega} = T^{-1} \sum_{t=1}^T \hat{u}_t \hat{u}'_t$ as before and, hence, $\hat{\Omega}$ is the residual covariance estimator from the restricted auxiliary model (4.4). Because the scaling matrix used in (4.5) is a consistent estimator of $\Sigma_{U \cdot X} \otimes \Omega$, it follows immediately from Proposition 1 that

$$Q_{BG}^* \xrightarrow{d} \chi^2(hK^2).$$

Several remarks are worth making regarding this result.

Remark 1. Notice that the regressors $y'_{t-1}\xi_{\perp} \otimes \hat{\alpha}$ and $-\hat{\alpha}(t-1)$ may be deleted from the auxiliary model (4.4) without affecting the limiting distribution of the test. This result holds because the estimators of β and τ are asymptotically independent of the estimators of γ and ϕ . As far as the small sample properties of the test are concerned, it is not clear that deleting these terms is a good idea because the independence does not hold in finite samples. We will denote the statistic obtained from the auxiliary model without the regressors $y'_{t-1}\xi_{\perp} \otimes \hat{\alpha}$ and $-\hat{\alpha}(t-1)$ by Q_{BG} in the following. \square

Remark 2. As mentioned in the previous section, Proposition 1 holds also if there are, for example, seasonal dummies in the VECM (2.1). Because the limiting distribution of the LM test statistics follows directly from the proposition, it is clear that the LM tests can also be used to test for residual AC in models with seasonal dummies. In that case the seasonal dummies should also be included in the auxiliary model (4.4). Also, from Extension 3 of Proposition 1 it follows that Q_{BG}^* has the same limiting null distribution if the cointegrating rank is not taken into account and the model is estimated in unrestricted form. Equivalently, the corresponding levels version may be estimated. \square

Remark 3. There may be parameter restrictions on ϕ so that $\phi = \phi(\theta)$, as discussed in Extension 2 in the previous section. Again, the limiting distributions of the LM statistics are unaffected. Here the restrictions have to be taken into account in the auxiliary model in the usual way, however. \square

Remark 4. Rather than considering the full VAR(h) model (4.2) for the error term, a single lag as in $u_t = B_h u_{t-h} + \varepsilon_t$ may be used in the alternative. In that case the null hypothesis is $H_0 : B_h = 0$. The relevant test statistic is obtained by replacing \hat{U}_t with \hat{u}_{t-h} and considering an auxiliary model just like (4.4) or the corresponding version without the terms $y'_{t-1}\xi_{\perp} \otimes \hat{\alpha}$ and $-\hat{\alpha}(t-1)$. In that case the limiting distribution of the LM statistic is $\chi^2(K^2)$. This version of the test is, for example, given in EViews 4. \square

Remark 5. It can be shown that the Breusch-Godfrey statistic can be written alternatively as

$$Q_{BG} = T \left(K - \text{tr}(\hat{\Omega}^{-1} \hat{\Omega}_e) \right),$$

where $\hat{\Omega}_e$ is the unrestricted residual covariance estimator of the auxiliary model used for the test (see Edgerton & Shukur (1999)). Instead of using this LM version of the test statistic, the asymptotically equivalent likelihood ratio (LR) or Wald versions may be used. For example,

$$Q_{LR} = T(\log \det \hat{\Omega} - \log \det \hat{\Omega}_e) \quad (4.6)$$

or

$$Q_W = T \left(\text{tr}(\hat{\Omega}_e^{-1} \hat{\Omega}) - K \right)$$

may be considered. The corresponding statistics based on the auxiliary model (4.4) with regressors related to the scores of the cointegration parameters will be denoted by Q_{LR}^* and Q_W^* , respectively. \square

In the next subsection the portmanteau test will be discussed.

4.2 Portmanteau Test

Although portmanteau tests may also be regarded as tests for the pair of hypotheses in (4.3), it may be more natural to think of them as tests for the null hypothesis

$$H_0 : E(u_t u'_{t-i}) = 0, \quad i = 1, 2, \dots, \quad (4.7)$$

which is tested against the alternative that at least one of these autocovariances and, hence, one autocorrelation is nonzero. The basic form of the portmanteau test statistic is

$$Q_P = T \sum_{j=1}^h \text{tr}(\hat{C}'_j \hat{\Omega}^{-1} \hat{C}_j \hat{\Omega}^{-1}) = T \hat{\mathbf{c}}' (I_h \otimes \hat{\Omega} \otimes \hat{\Omega})^{-1} \hat{\mathbf{c}}. \quad (4.8)$$

Appealing to Lemma 1, the limiting distribution of this statistic can be derived by using the auxiliary regression model (2.4), with β treated as known. Transform x_t to

$$z_t = \xi \beta' x_t + \beta_{\perp} \xi'_{\perp} \Delta x_t$$

and define the lag polynomial matrix

$$D(L) = [\Gamma(L) \xi \Delta - \alpha L : \Gamma(L) \beta_{\perp}] [\beta : \xi_{\perp}]',$$

where $\Gamma(L) = I_K - \sum_{j=1}^{p-1} \Gamma_j L^j$. Note that $D(L)$ is of order p with $D(L) = I_K - \sum_{j=1}^p D_j L^j$. From Lemma 1 of Saikkonen (2003) and the subsequent discussion it follows that we can transform model (2.4) to

$$z_t = \sum_{j=1}^p D_j z_{t-j} + u_t. \quad (4.9)$$

Note that this transformation does not affect the error term and that the process z_t is stationary. Moreover, the parameter matrices D_1, \dots, D_p are smooth functions of the parameters $\alpha, \Gamma_1, \dots, \Gamma_{p-1}$ (see the definition of $D(L)$ and recall that β (and ξ) are treated as known). Using the LS estimators $\tilde{\alpha}, \tilde{\Gamma}_1, \dots, \tilde{\Gamma}_{p-1}$ of (2.4) one can obtain Gaussian ML estimators of D_1, \dots, D_p in an obvious way. The residuals \tilde{u}_t can then be considered as residuals from a Gaussian ML estimation of (4.9). Thus, we can conclude that the approximate distribution of the portmanteau test can be obtained by applying the general result of Ahn (1988),

$$Q_P \approx \chi^2(hK^2 - K^2(p-1) - Kr)$$

in large samples if h is also large. Notice that due to the restrictions in the D_j parameter matrices the degrees of freedom are different than for a stationary VAR(p) model where the degrees of freedom are $hK^2 - K^2p$ (e.g., Lütkepohl (1991)).

A related statistic with potentially superior small sample properties is the adjusted portmanteau statistic,

$$Q_P^* = T^2 \sum_{j=1}^h \frac{1}{T-j} \text{tr}(\hat{C}_j' \hat{C}_0^{-1} \hat{C}_j \hat{C}_0^{-1}),$$

(see, e.g., Hosking (1980)). Its asymptotic properties are the same as those of Q_P . A number of other small sample adjustments with equivalent asymptotic properties have also been considered in the literature for stationary models (e.g., Li & McLeod (1981), Johansen (1995, p. 21/22)). They could be used for the presently considered case of cointegrated processes as well.

If there are parameter restrictions for $\alpha, \Gamma_1, \dots, \Gamma_{p-1}$ the limiting theory still holds. However, the test statistics have approximate $\chi^2(df)$ distributions, where the degrees of freedom $df = hK^2 - \#\{\text{freely estimated parameters in } \alpha \text{ and } \Gamma_j, j = 1, \dots, p-1\}$.

5 Simulations

We check the small sample properties of the residual AC tests by using Monte Carlo simulations because earlier studies for stationary processes (see Doornik (1996) and Edgerton & Shukur (1999)) have shown that some tests have poor small sample properties and relying on their asymptotic properties may be quite misleading. We therefore investigate the small sample properties of different test variants discussed in Section 4 in the context of VECMs. In particular, we consider the two variants of the LM tests, Q_{BG}^* and Q_{BG} , as well as the related LR and Wald versions of the BG test in our Monte Carlo study and compare their properties. Moreover, the portmanteau tests, Q_P and Q_P^* from Section 4.2 are considered. The objective of our simulations is to see how the test versions considered in the previous section perform in small samples when they are applied to cointegrated processes. It is not our intention to provide a full scale investigation of all potential modifications of the statistics that have been proposed for stationary processes.

5.1 Monte Carlo Design

Our Monte Carlo experiments are based on the following DGP specification

$$\Delta y_t = \nu + \alpha(\beta' y_{t-1} - \tau(t-1)) + \Gamma_1 \Delta y_{t-1} + u_t. \quad (5.1)$$

Using this DGP structure a number of experiments have been conducted varying factors that are likely to influence the small sample performance of different AC tests. In particular, we change the dimension of the system ($K = 2, \dots, 5$), the cointegration rank ($r = 1, \dots, K-1$), the parameters of the loading coefficients α , the dynamic structure Γ_1 and the structure of the error covariance matrix. Instead of reporting all Monte Carlo results we present a few typical results in Section 5.2. Most of the results will be illustrated using a variant of the 3-dimensional DGP (a) from Table 1. The cointegration matrix β of this DGP contains typical $(1, -1)$ relations, which are loaded into some of the VECM equations by the parameters a_1 . The dynamic correlation structure given by Γ_1 can be controlled by setting γ and γ_1 as in Edgerton & Shukur (1999). Trend term and intercept parameters are set by $\bar{\tau}$ and $\bar{\nu}$, respectively. Cross-equation error correlation is given by ρ , whereas the AC structure of the error u_t can be controlled by the parameters π_1 and π_2 . The effects of increasing the

dimension is illustrated by using the 5-dimensional DGP (b) from Table 1. We give the actual parameter values for different experiments in Table 2.

For each DGP variant given in Table 2 we have simulated $M = 1000$ sets of time series data for y_t using the levels version of (5.1) such that $T = 50, 100, 200, 500, 1000$ observations can be used for estimation of the VECM model. We use the Johansen RR regression on a VAR with correct lag length $p = 2$ and obtain an estimate of β and τ using the correct rank restriction r . This estimate is used to setup a VECM for obtaining the residuals used in AC testing. In each Monte Carlo replication we have applied the Q_{BG}, Q_{LR} , and Q_W tests, the variants with cointegration terms ($y'_{t-1}\xi_{\perp} \otimes \hat{\alpha}$ and $-\hat{\alpha}(t-1)$) in the auxiliary regression (Q_{BG}^*, Q_{LR}^* , and Q_W^*) and both variants of the portmanteau test (Q_P and Q_P^*) to the residuals of the VECM. To investigate the size and power properties of the tests we have recorded the relative rejection frequencies for different values of h . Results of the Monte Carlo study will be discussed in the following subsection.

5.2 Monte Carlo Results

Table 3 lists the empirical sizes of different test variants for the three-dimensional DGP (a) with $h = 1, 5, 10, 15$. The nominal size in all experiments is 0.05. The results have been obtained by using the parameter variant (1) given in Table 2. Note that no portmanteau test results are given for $h = 1$ because these tests are only valid for h larger than the lag length. Moreover, no results for the BG tests are given when $T = 50$ and $h = 15$ because for this case the number of parameters in the auxiliary model is too large relative to the number of observations available.

The first obvious result from Table 3 is related to the relative performance of the BG test variants. In contrast to Q_{BG} , the Q_{LR} and Q_W variants are both severely oversized in small samples, which is especially apparent from results for $h > 5$. This result is robust across all our Monte Carlo experiments, which indicates that the LM variant of the BG test is preferable to the LR and Wald versions in applied work. A similar result was also found by Edgerton & Shukur (1999) for stationary processes and, in a related context, by Mizon & Hendry (1980). It may be worth noting that small sample corrections are available for the LR and Wald tests for stationary processes (see Edgerton & Shukur (1999)) which may improve the performance of these tests also for VECMs. We do not consider such modifications here

because Edgerton & Shukur (1999) find that they only partially correct the size distortions in stationary systems.

Comparing results for Q_{BG} with those for Q_{BG}^* reveals that including $(y'_{t-1}\xi_{\perp} \otimes \hat{\alpha})$ and $-\hat{\alpha}(t-1)$ in the auxiliary regression is not advantageous in small sample situations. On the contrary, the variant without these additional terms typically outperforms Q_{BG}^* in the sense that its size is closer to the nominal size of the test. With the exception of $T = 50$, the Q_{BG} variant has a size reasonably close to the nominal level of 5%.

Not surprisingly, the portmanteau test variants are oversized for $h = 5$, emphasizing that they are only valid when h is considerably larger than the lag length of the model. Accordingly, the size decreases when lags up to $h = 10$ or $h = 15$ are used. When using a sufficiently large number of lags, we find the standard portmanteau statistic Q_P to have an empirical size smaller than the nominal one (see Panels (c) and (d) of Table 3) in small samples which is in line with earlier results for stationary processes (e.g., Hosking (1980)) and was the reason for proposing the adjusted statistic in the first place. In contrast, the modified statistic Q_P^* typically has size close to the nominal level for large h . Results from Panels (b) through (d) suggest that the size of Q_P^* decreases with increasing h . Therefore, we also checked results for Q_P^* with $h = 20$ and found that the size did not drop significantly below the nominal level. Overall, the results for the portmanteau tests suggest that Q_P^* works better than Q_P in models with sample sizes common in applied work.

All results in Table 3 have been obtained by imposing the correct cointegration rank. In empirical work, however, the absence of residual AC in the initial VAR model is often tested before the cointegration analysis. Therefore, we also checked the performance of the AC tests when they are applied to the residuals of an unrestricted VAR in levels and give results in Table 4. In this situation, of course, including the BG test variant with terms related to cointegration would not make sense and hence, only one variant of each test is considered here. Again we find that Q_{BG} outperforms the LR and the Wald variant. Moreover, we find that using the test on the unrestricted VAR increases the size substantially. Comparing, e.g., Panels (a) from Table 3 and Table 4 shows that, for $T = 100$, the size of Q_{BG} increases from 0.054 in the VECM to 0.076 in the unrestricted VAR. For the portmanteau tests we computed the degrees of freedom in two different ways. First, they are computed as in the VECM assuming the correct cointegration rank (denoted as Q_P and Q_P^*). In this case, the results are

similar to estimating the VECM. Second, the degrees of freedom are calculated based on the usual formula for unrestricted VAR models (denoted as Q_P^{VAR-DF} and $Q_P^{*,VAR-DF}$). Doing so almost doubles the size of the portmanteau tests. In other words, using the information on the cointegration properties is important for the portmanteau tests. As a consequence, an uncritical application of the portmanteau tests to a levels VAR is problematic if there are integrated and possibly cointegrated variables. In that case, a VECM should be specified and then the tests should be applied or the BG tests should be used.

Table 5 shows results for a large-dimensional system with 5 variables. While the relative performance of the tests is unchanged, the size properties are no longer satisfactory, typically exceeding 10% in small samples (e.g. for $T = 100$). Consequently, using the AC tests in large dimensional systems requires some extra care as the true null hypothesis is rejected too often in this case.

In addition to the foregoing results, we have also obtained results for different parameter variants of our DGPs. In general, a similar picture of the relative performance of tests is obtained by varying the properties of the underlying DGP. Introducing cross-equation error correlation (variant (2) from Table 2), changing the cointegration structure (variant (3)) and changing the dynamic correlation structure (variant (4)) does not affect the general results. Therefore, the detailed results are not reported here. We have also varied the cointegration rank of the DGPs and find again a similar picture.

A final set of Monte Carlo experiments has been conducted to investigate the power properties of AC tests. As a typical result we give the relative rejection frequencies of Q_{BG} , Q_P and Q_P^* at $h = 5$ and $h = 10$ for different values of π_1 . Hence, the actual process has an autoregressive error structure. Figure 1 shows the power for the 3-dimensional DGP with parameters according to variant (5) from Table 2. For $T = 50$ and $h = 5$ the modified portmanteau test seems to be more powerful than the BG tests at a first sight. However, most of the difference can be attributed to the difference in size (see Panel (b) of the Table 3). Accordingly, there is almost no difference in the power lines of these two tests in Panel (b) of the Figure, where Q_{BG} and Q_P^* have about the same size. Moreover, the line for Q_P in this panel indicates that its low size is associated with a loss in power against alternatives. Increasing the sample size generally moves the power lines closer together. Again, the remaining differences are small and may be attributed to differences in the size of

different tests (see Table 3). Using a moving average process (variant (6)) as an alternative error process in the power simulation leads to similar results and therefore the corresponding graph is not shown here.

Overall, the results suggest some guidelines for practical work: In VECMs with typical dimension and sample sizes for macroeconomic time series analysis, the LM test variant without additional cointegration terms in the auxiliary regression has the most favorable size properties when the number of lags to be tested, h , is relatively small. For large h , degrees of freedom limitations may prohibit its use. In that situation the modified portmanteau test Q_P^* can be used. If h is large enough, Q_P^* has a size close to the nominal one if the degrees of freedom are computed according to the cointegration rank of the system. If no information on the cointegration rank is available and the cointegrated VAR model is estimated in levels, the LM version of the BG tests is better suited than the portmanteau test. The reason for this is that the degrees of freedom in the latter test vary according to the cointegration rank of the system. In addition, applied time series econometricians should keep in mind that all considered tests have a tendency to reject the true null too often in fairly large-dimensional systems and small samples. Therefore, some type of correction would be desirable in these cases. Some of the many small sample corrections proposed for stationary processes could be considered, for example. Alternatively, a bootstrap may help to correct the small sample distortions. Such extensions are beyond the scope of this study, however.

6 Conclusions

In this study we have investigated the asymptotic properties of residual autocovariances and ACs and standard tests for residual AC for VAR processes with cointegrated variables. The main results that emerge from the asymptotic analysis are as follows: (1) The asymptotic distributions of the residual autocovariances and autocorrelations can be derived as in the stationary case by assuming that the cointegration relations are known. (2) The Breusch-Godfrey LM test for residual AC based on an auxiliary VECM analogously to the stationary VAR case, has the same asymptotic χ^2 null distribution as in the stationary case. (3) The portmanteau statistic for residual AC has a different approximate χ^2 distribution than for stationary VARs. This result also implies that applying the usual (stationary) portmanteau

test for checking the residuals of a VAR model with integrated and cointegrated variables has no sound theoretical basis.

We have also performed a small Monte Carlo study to check the finite sample performance of the tests and found generally similar results as in earlier studies for stationary processes. For relatively small systems and moderately large samples, the Breusch-Godfrey Q_{BG} statistic can be recommended if only low order AC is tested. In contrast, the modified portmanteau test Q_p^* performs reasonably well if higher order AC is to be tested. For the performance of the latter test it is important, however, that the cointegration rank of the system is taken into account when the degrees of freedom for the approximate distribution are determined. The test tends to reject far too often if it is applied to VARs in levels of cointegrated variables without such adjustments. Also for higher dimensional systems, the asymptotic distribution is generally a poor guide for small sample inference based on these tests. Thus, one may consider small sample adjustments or perhaps bootstrap procedures to correct for the size distortions in these cases.

Our results suggest that some commercial software for VAR and VECM analysis should be modified. In particular, the p -values for the portmanteau tests reported by EViews are apparently based on an incorrect approximate distribution when VARs with cointegrated variables in levels or VECMs are considered. Moreover, asymptotic p -values could be provided by PcGive for tests of residual AC for VECMs based on our theoretical results. In contrast, proper tests for residual AC of VECMs are available in JMulTi (Lütkepohl & Krätzig (2004)).

Appendix. Proofs

In this Appendix the notation from the previous sections is used.

A.1 Proof of Lemma 1

As discussed after Proposition 1, we can use model (3.4). Hence,

$$\hat{u}_t = \Delta x_t - (X_t' \otimes I_K) \hat{\phi} - \hat{\alpha}(\hat{\beta}^* - \beta^*)' x_{t-1}^* - \hat{\nu}^{(0)}, \quad t = 1, \dots, T, \quad (A.1)$$

where $\hat{\phi} = \text{vec}[\hat{\alpha} : \hat{\Gamma}_1 : \dots : \hat{\Gamma}_{p-1}]$, $\hat{\beta}^{*'} = [\hat{\beta}' : \hat{\tau}^{(0)}]$, $\beta^{*'} = [\beta' : \tau^{(0)}]$ and $x_{t-1}^* = [x'_{t-1} : -(t-1)]'$. Denoting the LS estimator of ϕ from the model

$$\Delta x_t = (X_t' \otimes I_K)\phi + u_t \quad (A.2)$$

by $\tilde{\phi}$, we can express \hat{u}_t in (A.1) as

$$\hat{u}_t = \Delta x_t - (X_t' \otimes I_K)\tilde{\phi} + w_t = \tilde{u}_t + w_t, \quad (A.3)$$

where $w_t = (X_t' \otimes I_K)(\tilde{\phi} - \hat{\phi}) - \hat{\alpha}(\hat{\beta}^* - \beta^*)'x_{t-1}^* - \hat{\nu}^{(0)}$. Thus,

$$\hat{C}_j - \tilde{C}_j = T^{-1} \sum_{t=j+1}^T \tilde{u}_t w'_{t-j} + T^{-1} \sum_{t=j+1}^T w_t \tilde{u}'_{t-j} + T^{-1} \sum_{t=j+1}^T w_t w'_{t-j}, \quad (A.4)$$

and it suffices to show that each of the three terms on the right-hand side (r.h.s.) is of order $O_p(T^{-1})$. To this end we shall first demonstrate that

$$\hat{\phi} - \tilde{\phi} = O_p(T^{-1}). \quad (A.5)$$

Define $\hat{X}_t^* = [(\hat{\beta}^{*'} x_{t-1}^*)', \Delta x'_{t-1}, \dots, \Delta x'_{t-p+1}]'$. Then $\hat{\phi}$ and $\hat{\nu}^{(0)}$ can be obtained by LS estimation from the regression model

$$\Delta x_t = (\hat{X}_t^{*'} \otimes I_K)\phi + \nu^{(0)} + u_t^*, \quad (A.6)$$

where $u_t^* = u_t - \alpha(\hat{\beta}^* - \beta^*)'x_{t-1}^*$. Using (3.5), the fact that $\tau^{(0)} = 0$ and well-known properties of stationary and I(1) processes (see Johansen (1995, Appendix B.7)), it is straightforward to check that the sample means of \hat{X}_t^* , u_t^* and u_t are of order $O_p(T^{-1/2})$. From this it further follows, using standard LS formulas, that

$$\hat{\phi} = \phi + \left(\left(\sum_{t=1}^T \hat{X}_t^* \hat{X}_t^{*'} \right)^{-1} \otimes I_K \right) \sum_{t=1}^T (\hat{X}_t^* \otimes I_K) u_t^* + O_p(T^{-1}). \quad (A.7)$$

Defining $X_t^* = [(\beta^{*'} x_{t-1}^*)', \Delta x'_{t-1}, \dots, \Delta x'_{t-p+1}]'$, we get $\hat{X}_t^* - X_t^* = [((\hat{\beta}^* - \beta^*)' x_{t-1}^*)' : 0]'$. Hence, it follows from standard limit results for stationary and I(1) processes that

$$T^{-1} \sum_{t=1}^T (\hat{X}_t^* \otimes I_K) u_t^* = T^{-1} \sum_{t=1}^T (X_t^* \otimes I_K) u_t + O_p(T^{-1})$$

and

$$T^{-1} \sum_{t=1}^T \hat{X}_t^* \hat{X}_t^{*'} = T^{-1} \sum_{t=1}^T X_t^* X_t^{*'} + O_p(T^{-1}).$$

Using these results and

$$\tilde{\phi} = \phi + \left(\left(\sum_{t=1}^T X_t X_t' \right)^{-1} \otimes I_K \right) \sum_{t=1}^T (X_t \otimes I_K) u_t \quad (A.8)$$

together with (A.7), it is finally straightforward to obtain (A.5).

Now consider the first term on the r.h.s. of (A.4). By the definition of w_t ,

$$T^{-1} \sum_{t=j+1}^T \tilde{u}_t w_{t-j}' = T^{-1} \sum_{t=j+1}^T \tilde{u}_t (\hat{\phi} - \tilde{\phi})' (X_t \otimes I_K) - T^{-1} \sum_{t=j+1}^T \tilde{u}_t x_t^{*'} (\hat{\beta}^* - \beta^*) \hat{\alpha}' - T^{-1} \sum_{t=j+1}^T \tilde{u}_t \hat{\nu}^{(0)'}$$

Because \tilde{u}_t is the LS residual from a standard stationary regression model, limit results for stationary and I(1) processes already used along with (3.5) and (A.5) show that the first two terms on the r.h.s. are of order $O_p(T^{-1})$. That the same is true for the last term follows by observing that the sample mean of \tilde{u}_t is of order $O_p(T^{-1/2})$ and $\hat{\nu}^{(0)} = O_p(T^{-1/2})$ because $\nu^{(0)} = 0$. Thus, the desired result has been established for the first term on the r.h.s. of (A.4) and similar arguments clearly apply for the second term. The third term involves no new features either, so details are omitted.

A.2 Proof of Proposition 1

According to Lemma 1, it suffices to show the proposition for the \tilde{C}_j . We therefore consider the regression model (A.2) and the LS estimator $\tilde{\phi}$. Defining $\tilde{U}_t = [\tilde{u}_{t-1}', \dots, \tilde{u}_{t-h}']'$ gives

$$\tilde{\mathbf{c}} = \text{vec}([\tilde{C}_1, \dots, \tilde{C}_h]) = T^{-1} \sum_{t=1}^T (\tilde{U}_t \otimes I_K) \tilde{u}_t,$$

where $\tilde{u}_t = 0$ for $t \leq 0$. Let \mathbf{c} be the counterpart of $\tilde{\mathbf{c}}$ obtained by replacing \tilde{u}_t by u_t . Because X_t is stationary and $E(X_t u_{t+j}') = 0$, $j \geq 0$, it is straightforward to check that

$$\begin{aligned} \tilde{\mathbf{c}} &= \mathbf{c} + T^{-1} \sum_{t=1}^T (U_t \otimes I_K) (\tilde{u}_t - u_t) + O_p(T^{-1}) \\ &= \mathbf{c} - T^{-1} \sum_{t=1}^T (U_t X_t' \otimes I_K) (\tilde{\phi} - \phi) + O_p(T^{-1}) \end{aligned}$$

(cf. Ahn (1988)). Denote $M_{UX} = T^{-1} \sum_{t=1}^T U_t X_t' = M_{XU}'$ and define M_{UU} and M_{XX} analogously. From the preceding representation of \mathbf{c} and (A.8) we then find that

$$T^{1/2} \tilde{\mathbf{c}} = [I_{hK^2} : -(M_{UX} M_{XX}^{-1} \otimes I_K)] T^{-1/2} \sum_{t=1}^T \begin{bmatrix} U_t \otimes I_K \\ X_t \otimes I_K \end{bmatrix} u_t + O_p(T^{-1/2}).$$

Proposition 1 follows from this result, $\text{plim } T^{-1} \sum_{t=1}^T u_t u_t' = \Omega$ and

$$\frac{1}{\sqrt{T}} \left(\sum_{t=1}^T \begin{bmatrix} U_t \otimes I_K \\ X_t \otimes I_K \end{bmatrix} u_t \right) \xrightarrow{d} N(0, \bar{\Sigma}), \quad (A.9)$$

where

$$\bar{\Sigma} = \begin{bmatrix} \Sigma_{UU} & \Sigma_{UX} \\ \Sigma_{XU} & \Sigma_{XX} \end{bmatrix} \otimes \Omega$$

with $\Sigma_{UU} = I_h \otimes \Omega$. The limit result in (A.9) follows from a standard martingale central limit theorem using (2.3) (see Johansen (1995, Appendix B.7)).

References

- Ahn, S. K. (1988). Distribution for residual autocovariances in multivariate autoregressive models with structured parameterization, *Biometrika* **75**: 590–593.
- Doornik, J. A. (1996). Testing vector error autocorrelation and heteroscedasticity, unpublished paper, Nuffield College.
- Edgerton, D. & Shukur, G. (1999). Testing autocorrelation in a system perspective, *Econometric Reviews* **18**: 343–386.
- Hosking, J. R. M. (1980). The multivariate portmanteau statistic, *Journal of the American Statistical Association* **75**: 602–608.
- Hosking, J. R. M. (1981a). Equivalent forms of the multivariate portmanteau statistic, *Journal of the Royal Statistical Society* **B43**: 261–262.
- Hosking, J. R. M. (1981b). Lagrange-multiplier tests of multivariate time series models, *Journal of the Royal Statistical Society* **B43**: 219–230.
- Johansen, S. (1991). Estimation and hypothesis testing of cointegration vectors in Gaussian vector autoregressive models, *Econometrica* **59**: 1551–1581.
- Johansen, S. (1995). *Likelihood-based Inference in Cointegrated Vector Autoregressive Models*, Oxford University Press, Oxford.

- Li, W. K. & McLeod, A. I. (1981). Distribution of the residual autocorrelations in multivariate ARMA time series models, *Journal of the Royal Statistical Society* **B43**: 231–239.
- Lütkepohl, H. (1991). *Introduction to Multiple Time Series Analysis*, Springer Verlag, Berlin.
- Lütkepohl, H. & Krätzig, M. (eds) (2004). *Applied Time Series Econometrics*, Cambridge University Press, Cambridge.
- Mizon, G. E. & Hendry, D. F. (1980). An empirical application and Monte Carlo analysis of tests of dynamic specification, *Review of Economic Studies* **47**: 21–45.
- Reinsel, G. C. & Ahn, S. K. (1992). Vector autoregressive models with unit roots and reduced rank structure: Estimation, likelihood ratio test, and forecasting, *Journal of Time Series Analysis* **13**: 353–375.
- Saikkonen, P. (2001a). Consistent estimation in cointegrated vector autoregressive models with nonlinear time trends in cointegrating relations, *Econometric Theory* **17**: 296–326.
- Saikkonen, P. (2001b). Statistical inference in cointegrated vector autoregressive models with nonlinear time trends in cointegrating relations, *Econometric Theory* **17**: 327–356.
- Saikkonen, P. (2003). Stability results for nonlinear error correction models, *Technical report*, University of Helsinki.

Table 1: DGPs Used in Monte Carlo Experiment

| | | |
|---|---|---|
| (a) $K = 3, r = 2:$ | $\alpha = \begin{pmatrix} a_1 & 0 \\ 0 & a_1 \\ 0 & 0 \end{pmatrix},$ | $\beta' = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}$ |
| | | |
| (b) $K = 5, r = 2:$ | $\alpha = \begin{pmatrix} a_1 & 0 \\ 0 & a_1 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix},$ | $\beta' = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{pmatrix}$ |
| | | |
| $\Gamma_1 = \Gamma^* \quad \text{with} \quad \gamma_{ij}^* = \begin{cases} \gamma & j = i \\ \gamma_1 & j = i + 1 \text{ or } i, j = K, 1 \\ 0 & \text{otherwise} \end{cases} \quad \tau = \bar{\tau} \mathbf{1}_r, \quad \nu = \bar{\nu} \mathbf{1}_K$ | | |
| $u_t = B_1 u_{t-1} + e_t + A_1 e_{t-1}, \quad B_1 = \pi_1 I_K, \quad A_1 = \pi_2 I_K, \quad e_t \sim \text{iid } N(0, \Omega_e)$ | | |
| $\Omega_e = \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 & \dots & \rho\sigma_1\sigma_K \\ \rho\sigma_2\sigma_1 & \sigma_2^2 & & \vdots \\ \vdots & & \ddots & \rho\sigma_{K-1}\sigma_K \\ \rho\sigma_K\sigma_1 & \dots & \rho\sigma_K\sigma_{K-1} & \sigma_K^2 \end{pmatrix}$ | | |

Note: $\mathbf{1}_K$ denotes a $(K \times 1)$ vector of ones.

Table 2: Parameter Values of DGPs

| | Experiment | | | | | |
|--------------|------------|------|------|------|-------------|-------------|
| | (1) | (2) | (3) | (4) | (5) | (6) |
| a_1 | -0.2 | -0.2 | -0.1 | -0.2 | -0.2 | -0.2 |
| γ | 0.5 | 0.5 | 0.5 | 0.7 | 0.5 | 0.5 |
| γ_1 | -0.2 | -0.2 | -0.2 | -0.1 | -0.2 | -0.2 |
| $\bar{\tau}$ | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| $\bar{\nu}$ | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| σ_i | 1 | 1 | 1 | 1 | 1 | 1 |
| ρ | 0 | 0.7 | 0 | 0.7 | 0 | 0 |
| π_1 | 0 | 0 | 0 | 0 | 0, ..., 0.8 | 0 |
| π_1 | 0 | 0 | 0 | 0 | 0 | 0, ..., 0.8 |

Table 3: Empirical Sizes of AC Tests, DGP (a): $K = 3$, $r = 2$, variant (1)

| (a) $h = 1$ | | | | | | (b) $h = 5$ | | | | |
|--------------|-------|-------|-------|-------|-------|--------------|-------|-------|-------|-------|
| T | 50 | 100 | 200 | 500 | 1000 | 50 | 100 | 200 | 500 | 1000 |
| Q_{BG} | 0.111 | 0.054 | 0.042 | 0.041 | 0.042 | 0.126 | 0.079 | 0.056 | 0.056 | 0.049 |
| Q_{LR} | 0.169 | 0.073 | 0.052 | 0.043 | 0.044 | 0.651 | 0.251 | 0.101 | 0.071 | 0.055 |
| Q_W | 0.221 | 0.091 | 0.058 | 0.043 | 0.045 | 0.911 | 0.415 | 0.186 | 0.094 | 0.062 |
| Q_{BG}^* | 0.135 | 0.062 | 0.046 | 0.042 | 0.043 | 0.228 | 0.092 | 0.061 | 0.059 | 0.050 |
| Q_{LR}^* | 0.187 | 0.079 | 0.054 | 0.043 | 0.044 | 0.768 | 0.274 | 0.113 | 0.077 | 0.056 |
| Q_W^* | 0.246 | 0.102 | 0.060 | 0.043 | 0.045 | 0.954 | 0.456 | 0.195 | 0.097 | 0.064 |
| Q_P | | | | | | 0.104 | 0.094 | 0.089 | 0.109 | 0.092 |
| Q_P^* | | | | | | 0.176 | 0.134 | 0.104 | 0.115 | 0.093 |
| (c) $h = 10$ | | | | | | (d) $h = 15$ | | | | |
| T | 50 | 100 | 200 | 500 | 1000 | 50 | 100 | 200 | 500 | 1000 |
| Q_{BG} | 0.116 | 0.048 | 0.050 | 0.057 | 0.050 | | 0.038 | 0.038 | 0.053 | 0.051 |
| Q_{LR} | 1.000 | 0.558 | 0.205 | 0.091 | 0.067 | | 0.919 | 0.380 | 0.138 | 0.091 |
| Q_W | 1.000 | 0.892 | 0.438 | 0.149 | 0.092 | | 1.000 | 0.781 | 0.249 | 0.142 |
| Q_{BG}^* | 0.370 | 0.079 | 0.054 | 0.060 | 0.050 | | 0.077 | 0.052 | 0.059 | 0.052 |
| Q_{LR}^* | 1.000 | 0.630 | 0.227 | 0.095 | 0.067 | | 0.951 | 0.430 | 0.146 | 0.095 |
| Q_W^* | 1.000 | 0.922 | 0.464 | 0.157 | 0.092 | | 1.000 | 0.803 | 0.256 | 0.144 |
| Q_P | 0.022 | 0.020 | 0.032 | 0.051 | 0.052 | 0.005 | 0.010 | 0.018 | 0.044 | 0.048 |
| Q_P^* | 0.114 | 0.069 | 0.051 | 0.058 | 0.058 | 0.095 | 0.052 | 0.041 | 0.050 | 0.060 |

Note: The nominal size of all tests is 5%.

Table 4: Empirical Sizes of AC Tests Based on VAR in Levels, DGP (a): $K = 3$, $r = 2$, variant (1)

| (a) $h = 1$ | | | | | | (b) $h = 5$ | | | | |
|------------------|-------|-------|-------|-------|-------|-------------|-------|-------|-------|-------|
| T | 50 | 100 | 200 | 500 | 1000 | 50 | 100 | 200 | 500 | 1000 |
| Q_{BG} | 0.149 | 0.076 | 0.048 | 0.043 | 0.042 | 0.324 | 0.111 | 0.063 | 0.058 | 0.054 |
| Q_{LR} | 0.214 | 0.097 | 0.057 | 0.044 | 0.042 | 0.851 | 0.318 | 0.126 | 0.076 | 0.059 |
| Q_W | 0.282 | 0.113 | 0.066 | 0.046 | 0.043 | 0.977 | 0.487 | 0.201 | 0.102 | 0.064 |
| Q_P | | | | | | 0.111 | 0.091 | 0.086 | 0.107 | 0.092 |
| Q_P^* | | | | | | 0.189 | 0.121 | 0.099 | 0.110 | 0.093 |
| Q_P^{VAR-DF} | | | | | | 0.212 | 0.182 | 0.181 | 0.189 | 0.177 |
| $Q_P^{*,VAR-DF}$ | | | | | | 0.331 | 0.229 | 0.203 | 0.197 | 0.185 |

Note: The nominal size of all tests is 5%.

Table 5: Empirical Sizes of AC Tests, DGP (b): $K = 5$, $r = 2$, variant (1)

| (a) $h = 5$ | | | | | | (b) $h = 10$ | | | | |
|-------------|-------|-------|-------|-------|-------|--------------|-------|-------|-------|-------|
| T | 50 | 100 | 200 | 500 | 1000 | 50 | 100 | 200 | 500 | 1000 |
| Q_{BG} | 0.454 | 0.135 | 0.086 | 0.063 | 0.054 | | 0.152 | 0.083 | 0.061 | 0.052 |
| Q_{LR} | 1.000 | 0.719 | 0.284 | 0.110 | 0.070 | | 1.000 | 0.728 | 0.215 | 0.105 |
| Q_W | 1.000 | 0.965 | 0.535 | 0.187 | 0.095 | | 1.000 | 0.988 | 0.463 | 0.194 |
| Q_{BG}^* | 0.815 | 0.220 | 0.102 | 0.068 | 0.055 | | 0.353 | 0.123 | 0.068 | 0.055 |
| Q_{LR}^* | 1.000 | 0.819 | 0.336 | 0.124 | 0.075 | | 1.000 | 0.784 | 0.231 | 0.112 |
| Q_W^* | 1.000 | 0.979 | 0.594 | 0.197 | 0.098 | | 1.000 | 0.993 | 0.488 | 0.201 |
| Q_P | 0.176 | 0.150 | 0.140 | 0.136 | 0.100 | 0.015 | 0.028 | 0.037 | 0.054 | 0.055 |
| Q_P^* | 0.378 | 0.220 | 0.166 | 0.146 | 0.104 | 0.237 | 0.119 | 0.083 | 0.071 | 0.063 |

Note: The nominal size of all tests is 5%.




Figure 1: Powers of Breusch-Godfrey and portmanteau tests, DGP(a): $K = 3, r = 2$, variant (5).